



- Home
- About Us
- How to Help
- News & Events
- Disorders & Conditions
- Research Center



- » **For the Media**
- Expert Sources
  - Recent Press Releases
  - NARSAD News Kit
  - Research & Giving News

- » **Events**
- Galas & Scientific Symposia
  - Parlor Meetings & Seminars
  - Community Fundraisers

- » **Publications**
- NARSAD Research Quarterly
  - Free Brochures & Fact Sheets on Mental Illness
  - Annual Report

- » **Streaming Videos**
- Healthy Minds



**Research & Giving News Article** Jun. 3, 2009

EMAIL THIS PAGE
 PRINT THIS PAGE

### Using Light to Alter Neural Activity

Potential for Safe, Powerful New Tool for Neurological and Psychiatric Therapy

(Great Neck, N.Y. - May 26, 2009) — NARSAD Investigator Edward S. Boyden, Ph.D., and colleagues at the Massachusetts Institute of Technology, have demonstrated that optical methods can be used to manipulate neural activity in the nonhuman primate brain.

The research, reported in the journal *Neuron* on April 29, 2009, highlights a methodology for investigating the causal role of specific cell types and pathways in the nervous system; in neural computation, cognition and behavior. The findings, the authors stated, “open up the possibility of a new generation of ultraprecise neurological and psychiatric therapeutics via cell-type-specific optical neural control prosthetics” that may improve efficacy while reducing side effects.

“We anticipate that this work may lead to many novel therapies for blindness, Parkinson’s, epilepsy, and other disorders, said Dr. Boyden, the recipient of a 2008 NARSAD Young Investigator Award.

To understand how brain states and behaviors are generated by neural circuits, the researchers used a slow-acting virus, called a lentivirus, to target a light-activated protein channel, called channelrhodopsin-2 (ChR2), specific to excitatory neurons of the macaque frontal cortex.

Using a laser-coupled optical fiber in conjunction with a recording microelectrode, the team showed that activation of excitatory neurons resulted in well-timed excitatory and suppressive influences on neocortical neural networks. ChR2 was safely expressed, and could mediate optical neuromodulation, in primate neocortex over many months.

(This article was adapted with permission from *Neuron*.)

EMAIL THIS PAGE
 PRINT THIS PAGE

Media Contact

[Kristen Simone](#)  
516-829-0091, ext. 236

- Latest News from NARSAD
- Gene Linked to Autism Risk in Boys
  - Potential New Target for Depression Treatment
  - For Proper Brain Function, Cells Need to Keep the Beat
  - Using Light to Alter Neural Activity
  - Variant Gene May Impair Brain’s Communication in Schizophrenia and Bipolar

Spotlight

**Award-Winning “Healthy Minds” TV Series Presents Second Season**

[Click Here for Airtimes and Streaming Episodes](#)

Order these new books on schizophrenia and help NARSAD: [My Son’s Name Was Fred](#) by Gwill Newman, and [I Think I Scared Her](#) by Brooke Katz.



The HealthCentral Network  
[www.HealthCentral.com](http://www.HealthCentral.com)





**NARSAD** 60 Cutter Mill Road, Suite 404, Great Neck, New York 11021 USA    **phone** (800) 829-8289    **fax** (516) 487-6930    **email** [info@narsad.org](mailto:info@narsad.org)

[©NARSAD 2009](#) | [privacy policy](#) | [legal notices](#) | [disclaimers](#) | [sitemap](#) | [site help](#) | [contact us](#)